

SOLVING DEGENERACIES AT A β -BEAM BY COMBINING IONS

E. FERNANDEZ-MARTINEZ

I.F.T. and Dep. Física Teórica, Univ. Autonoma Madrid., E-28049, Madrid, Spain

We study how the eightfold-degeneracy in the (θ_{13}, δ) plane observed in $\gamma \sim 100$ β -beams can be reduced by periodically changing the ions in the storage ring. This “ions cocktail” allows to change the neutrino energy, at fixed γ , by choosing ions with different decay energies. We propose to combine the standard ${}^6\text{He}$ and ${}^{18}\text{Ne}$ beams with ${}^8\text{Li}$ and ${}^8\text{B}$ ones. These latter two ions have peaked $\nu_e \rightarrow \nu_\mu$ oscillation probabilities for $\gamma = 100$ at a baseline $L \sim 700$ Km. At this distance and this γ the oscillation probability of ${}^6\text{He}$ and ${}^{18}\text{Ne}$ neutrinos is at its second maximum. This setup is particularly suited for large enough values of θ_{13} (within reach at T2K-I) and it allows solving most of the eightfold-degeneracy, measuring θ_{13}, δ and the sign of the atmospheric mass difference for values of $\theta_{13} \geq 5^\circ$.

1 The Alternating Ions Scheme

The results of atmospheric, solar, accelerator and reactor neutrino experiments¹ show that flavour mixing occurs in the leptonic sector. Experiments have measured two mass differences, $\Delta m_{sol}^2 \approx 7.9 \times 10^{-5} \text{ eV}^2$ and $|\Delta m_{atm}^2| \approx 2.4 \times 10^{-3} \text{ eV}^2$. But only two out of the four parameters of the three-family leptonic mixing matrix U_{PMNS} are known: $\theta_{12} \approx 34^\circ$ and $\theta_{23} \approx 41.5^\circ$ ¹. The other two parameters, θ_{13} and δ , are still unknown: for θ_{13} direct searches at reactors give the upper bound $\theta_{13} \leq 11.5^\circ$, whereas for the leptonic CP-violating phase δ we have no information whatsoever. Two additional discrete unknowns are the sign of the atmospheric mass difference and the θ_{23} -octant. The two unknown parameters θ_{13} and δ can be measured in “appearance” experiments through $\nu_e \rightarrow \nu_\mu, \nu_\mu \rightarrow \nu_e$ (the “golden channel”²) and $\nu_e \rightarrow \nu_\tau$ (the “silver channel”³) oscillations. However, strong correlations between θ_{13} and δ and the presence of parametric degeneracies in the (θ_{13}, δ) parameter space,^{4–7} make the simultaneous measurement of the two variables extremely difficult. Most proposed solutions to these problems suggest the combination of different experiments and facilities, such as Super-Beam’s (of which T2K⁸ is the first approved one), β -beam’s⁹ or the Neutrino Factory¹⁰.

Here we propose to alleviate the parametric degeneracy in the (θ_{13}, δ) plane by combining β -beam’s obtained from the decay of several different ions¹¹. The β -beam concept⁹ involves producing a huge number of β -unstable ions, accelerating them to some reference energy, and letting them decay in the straight section of a storage ring, resulting in a very intense and pure ν_e or $\bar{\nu}_e$ beam. “Golden” sub-leading transitions, $\nu_e \rightarrow \nu_\mu$ and $\bar{\nu}_e \rightarrow \bar{\nu}_\mu$, can then be measured through muon observation in a distant detector.

In Tab. 1 we remind the relevant parameters for five ions: ${}^{18}\text{Ne}$, ${}^6\text{He}$, ${}^8\text{Li}$ and ${}^8\text{B}$. Consider first ${}^6\text{He}$ and ${}^{18}\text{Ne}$: as it was stressed in the literature, ${}^6\text{He}$ has the right half-life to be accelerated and stored such as to produce an intense $\bar{\nu}_e$ beam. This is also the case for ${}^{18}\text{Ne}$, the best candidate as β^+ -emitter. These two ions are the ones usually proposed as sources for the

Element	A/Z	$T_{1/2}$ (s)	Q_β eff (MeV)	Decay Fraction
^{18}Ne	1.8	1.67	3.41	92.1%
			2.37	7.7%
			1.71	0.2%
			13.92	100%
^8B	1.6	0.77		
^6He	3.0	0.81	3.51	100%
^8Li	2.7	0.83	12.96	100%

Table 1: A/Z , half-life and decay energy for two β^+ -emitters (^{18}Ne and ^8B) and two β^- -emitters (^6He and ^8Li).

neutrino beams.

The two ions we propose as an alternative to ^6He and ^{18}Ne as β^- and β^+ -emitters are ^8Li and ^8B , respectively. A novel way to produce intense beams of these ions was proposed in Ref. 12. ^8Li and ^8B have similar half-life, Z and A/Z to ^6He and ^{18}Ne , thus sharing the key characteristics needed for the bunch manipulation, but both ions have a much larger end-point energy than the two reference ions. As a consequence, for a fixed γ , a longer baseline is needed, and thus a smaller signal statistics is expected in the far detector. Therefore, the expected sensitivity to θ_{13} of such beams is smaller than that for a “standard” beam produced via ^6He and ^{18}Ne . However, when γ is limited (for example when using the CERN SPS as it is) it is then possible to reach higher neutrino energies using the same facility to accelerate the ions to be stored. If we combine different ions, we can (using the same facility and the same γ factor) produce neutrino beams of different L/E that can be used to disentangle many of the parametric degeneracies discussed before. As it was shown in the literature (see, for example, Refs. 15-17) degeneracies are indeed best lifted combining beams with different L/E .

Three classes of setups have been considered so far: $\gamma \sim 100$ (“low” γ), with a typical baseline of $O(100)$ Km^{13-14, 16-17}; $\gamma \sim 300$ (“medium” γ), with $L = O(700)$ Km^{13, 18-20}; and $\gamma \geq 1000$ (“high” γ), with baselines of several thousands kilometers,^{13, 19}.

We will compare results obtained with three different setups:

Standard “low” γ scenario $L = 130$ Km (CERN to Fréjus); $\gamma_{^6\text{He}} = \gamma_{^{18}\text{Ne}} = 120$. Both fluxes are tuned to be at the first oscillation peak. A given ion is accumulated in the storage ring for ten years.

Alternating ions “low” γ scenario $L = 650$ Km (CERN to Canfranc); $\gamma_{^8\text{Li}} = \gamma_{^8\text{B}} = 100$; $\gamma_{^6\text{He}} = \gamma_{^{18}\text{Ne}} = 120$. The ^8Li and ^8B fluxes are tuned at the first oscillation peak, whereas the ^6He and ^{18}Ne fluxes are tuned at the second oscillation peak. A given ion is accumulated in the storage ring for five years.

Standard “medium” γ scenario $L = 650$ Km (CERN to Canfranc); $\gamma_{^6\text{He}} = \gamma_{^{18}\text{Ne}} = 350$. Both fluxes are tuned to be at the first oscillation peak. A given ion is accumulated in the storage ring for ten years.

In all scenarios, a $\bar{\nu}_e$ flux of $2.9 \times 10^{18} \bar{\nu}_e$ per year or a ν_e flux of $1.1 \times 10^{18} \nu_e$ per year is aimed at the distant detector, regardless of the decaying ion. For a 5 year running time per each ion stored, the total luminosity is $1.45 \times 10^{19} \bar{\nu}_e$ and $5.5 \times 10^{18} \nu_e$ aimed at the far detector. We will consider a 1 Mton mass water Čerenkov detector (500 Kton fiducial mass), with a MEMPHYS design²¹. At the considered neutrino energies, this detector is believed to show a rather good neutrino energy reconstruction capability, allowing for a significant background rejection. Migration and background matrices for this detector have been taken from Ref. 18.




Figure 1: 90% C.L. contours for the standard “low” γ setup (left) and the alternating ions “low” γ setup (right). The input pair is labelled by a thick black square and corresponds to $\theta_{13} = 5^\circ, 8^\circ$ and $\delta = 45^\circ, -90^\circ$.




Figure 2: Comparison of the 3σ θ_{13} -sensitivity (left) and the $\text{sign}(\Delta m_{23}^2)$ -sensitivity (right) for the alternating ions “low” γ setup (solid) and the standard “medium” γ setup (dotted). The dashed line stands for the standard “low” γ setup in the left panel and a combination of “low” and “medium” γ in the right panel.

2 Solving degeneracies

In Fig. 1 we present our results for a fit to “experimental data” (generated as in Ref. 2) for the **standard “low” γ setup** (left plot) and the **alternating ions “low” γ setup** (right plot). The input pair is labelled by a thick black square. Lines represent 90% C.L. contours. The sign of Δm_{23}^2 has been chosen to be positive. Since the sign of Δm_{23}^2 and the θ_{23} -octant are unknown, fits to both $\text{sign}[\Delta m_{23}^2] = \pm 1$ and $\text{sign}[\tan(2\theta_{23})] = \pm 1$ have been performed (see Ref. 16 for a description of parametric degeneracies at the “low” γ β -beam).

As it can be seen, for every considered input pair the *Alternating Ions Scheme* reduces the eightfold-degeneracy in the (θ_{13}, δ) plane to a twofold-degeneracy. The so-called “intrinsic” degeneracy⁴ is always solved (as it is usual when combining information from neutrino beams with different L/E). Most importantly, the sign of the atmospheric mass difference is measured. This is not possible at the standard “low” γ setup since the baseline is too short to take advantage of matter effects to discriminate between hierarchical and inverted spectra.

In Fig. 2 we present our results for the θ_{13} -sensitivity (left panel) and $\text{sign}(\Delta m_{23}^2)$ -sensitivity (right panel) for the standard “low” γ setup (dashed lines in the left panel), the alternating ions “low” γ setup (solid lines) and the standard “medium” γ setup (dotted lines). Notice that sensitivities are defined as in Refs. 16 and 22, taking into account all of the parametric degeneracies. Lines represent 3σ contours. As it was expected, our *Alternating Ions Scheme* has the worst θ_{13} -sensitivity as a consequence of the lower statistics (Fig. 2, left). The decrease in statistics due to the longer baseline is compensated at the “medium” γ setup by an increase in

γ , providing a far better sensitivity than that of the standard “low” γ one, as noticed in Ref. 18. On the other hand (Fig. 2, right), the *Alternating Ions Scheme* can measure the sign of the atmospheric mass difference for $\sin^2(2\theta_{13}) \geq 0.04$ ($\theta_{13} \geq 5^\circ$). This must be compared with the standard “low” γ setup, with no sensitivity to $\text{sign}(\Delta m_{23}^2)$ due to its too short baseline. Even compared to the standard “medium” γ setup, with a much larger statistics, our scheme is particularly effective for small $|\delta|$, a region of the parameter space in which other setups are not working very well^a. This is a consequence of the combination of neutrino beams tuned at different oscillation peaks, something that guarantees that “sign clones” are located in different regions of the (θ_{13}, δ) parameter space also in the case of a vanishing δ . But this fact could also be exploited if higher γ is available. As we have discussed the “medium” γ setup clearly outperforms the lower γ ones in sensitivity to θ_{13} and to CP violation for small θ_{13} . Thus, if θ_{13} is not discovered at the next generation of experiments, such as T2K, the “medium” γ setup would be the best choice to measure θ_{13} and δ_{CP} . However, if nature has been unkind, we could find that we are in a region of the parameter space on Fig. 2 in which this setup has no sensitivity to the sign. An interesting possibility would then be to run the β -Beam at the lower γ , so that the oscillation probability is at its second peak for the same detector baseline, and combine it with the higher energy run at the first peak. This is depicted by the dashed line in the right panel of Fig. 2, where the regions in which the “medium” γ setup could not measure the sign are now also covered by the sensitivity curve.

In summary, we think that the *Alternating Ions Scheme* is particularly well-suited to solve degeneracies in the physics case where θ_{13} is relatively large (i.e. measurable at T2K-I). For smaller values of θ_{13} higher γ factors are required but the degeneracies could also be softened in this case by combining runs with different γ factors.

References

1. G. L. Fogli *et al.*, arXiv:hep-ph/0506083; G. L. Fogli *et al.*, arXiv:hep-ph/0506307.
2. A. Cervera *et al.*, Nucl. Phys. B **579** (2000) 17 [arXiv:hep-ph/0002108].
3. A. Donini *et al.*, Nucl. Phys. B **646** (2002) 321 [arXiv:hep-ph/0206034]; D. Autiero *et al.*, Eur. Phys. J. C **33** (2004) 243 [arXiv:hep-ph/0305185].
4. J. Burguet-Castell *et al.*, Nucl. Phys. B **608** (2001) 301 [arXiv:hep-ph/0103258].
5. H. Minakata and H. Nunokawa, JHEP **0110** (2001) 001 [arXiv:hep-ph/0108085].
6. G. L. Fogli and E. Lisi, Phys. Rev. D **54** (1996) 3667 [arXiv:hep-ph/9604415].
7. V. Barger *et al.*, Phys. Rev. D **65** (2002) 073023 [arXiv:hep-ph/0112119].
8. Y. Itow *et al.*, arXiv:hep-ex/0106019.
9. P. Zucchelli, Phys. Lett. B **532** (2002) 166; arXiv:hep-ex/0107006.
10. S. Geer, Phys. Rev. D **57** (1998) 6989 [Erratum-ibid. D **59** (1999) 039903] [arXiv:hep-ph/9712290]; A. De Rujula *et al.*, Nucl. Phys. B **547** (1999) 21 [arXiv:hep-ph/9811390].
11. A. Donini and E. Fernandez-Martinez, arXiv:hep-ph/0603261.
12. C. Rubbia *et al.*, arXiv:hep-ph/0602032.
13. J. Burguet-Castell *et al.*, Nucl. Phys. B **695**, 217 (2004) [arXiv:hep-ph/0312068].
14. J. Bouchez *et al.*, AIP Conf. Proc. **721** (2004) 37 [arXiv:hep-ex/0310059]; M. Mezzetto, J. Phys. G **29** (2003) 1771 [arXiv:hep-ex/0302007].
15. A. Donini *et al.*, JHEP **0406**, 011 (2004) [arXiv:hep-ph/0312072].
16. A. Donini *et al.*, Nucl. Phys. B **710** (2005) 402 [arXiv:hep-ph/0406132].
17. A. Donini *et al.*, Phys. Lett. B **621** (2005) 276 [arXiv:hep-ph/0411402].
18. J. Burguet-Castell *et al.*, Nucl. Phys. B **725**, 306 (2005) [arXiv:hep-ph/0503021].

^aThis was the case for the Neutrino Factory, also, see Ref. 22.

19. P. Huber *et al.*, Phys. Rev. D **73**, 053002 (2006) [arXiv:hep-ph/0506237]; F. Terranova *et al.*, Eur. Phys. J. C **38**, 69 (2004) [arXiv:hep-ph/0405081].
20. A. Donini *et al.*, arXiv:hep-ph/0604229.
21. J. E. Campagne *et al.*, arXiv:hep-ph/0603172.
22. A. Donini *et al.*, arXiv:hep-ph/0512038;